

DOI: 10.21767/2471-7975.100009

Hardcastle Takes a HIT! Commentary: Why Sprint Interval Training is Inappropriate for a Largely Sedentary Population

Sarah J Hardcastle^{1*} and Eduardo Caldas Costa²

¹Faculty of Health Sciences, Health Psychology and Behavioural Medicine Research Group, School of Psychology and Speech Pathology, Curtin University, Perth, WA, Australia

²Department of Physical Education, Federal University of Rio Grande do Norte, Natal, RN, Brazil

*Corresponding author: Sarah J. Hardcastle, Faculty of Health Sciences, Health Psychology and Behavioural Medicine Research Group, School of Psychology and Speech Pathology, Curtin University, Perth, WA, Australia, Tel: +61 8 9266 9266; E-mail: sarah.hardcastle@curtin.edu.au

Received date: November 23, 2016; Accepted Date: November 22, 2017; Published date: November 30, 2017

Citation: Hardcastle SJ, Costa EC (2017) Hardcastle Takes a HIT! Commentary: Why Sprint Interval Training is Inappropriate for a Largely Sedentary Population. Ann of Behav Sci Vol.3 No.3: 9.

Commentary

Our previous paper prompted a lively debate concerning the efficacy of sprint interval training (SIT) [1,2]. For clarification, SIT refers to protocols that involve supramaximal efforts (>100% maximal oxygen uptake) and high-intensity-interval-training (HIIT) involves 'vigorous' or 'near-maximal' efforts (target intensity: 80% to 100% peak heart-rate) [3]. In the current commentary, we return to the role of affect and perceived effort on exercise adherence raised in the recent commentary by Jung et al. [4]. We contend that the considerable anticipated effort required to participate in SIT and aversive psychological states experienced during such 'all out' supramaximal exercise may likely to damage adherence to such protocols.

Although we recognize the health-related physiological benefits of classic SIT, which involves up to six 30-s-sprints, we agree that 'considering the need for specialized equipment and the extremely elevated level of subject motivation, this form of training may not be safe, tolerable or practical for many individuals' [5]. Therefore, we contend that the considerable effort required to participate in classic SIT is such that it is unlikely to be experienced as pleasant and enjoyable for most people, especially those less active individuals [6].

More recently researchers have proposed SIT protocols with fewer and shorter sprints than classic SIT [7-9]. For example, Metcalfe et al. [7] have proposed the reduced-exertion high-intensity training, which involves two 'all out' 20-s-sprints, while Gillen et al. [8] have proposed a SIT protocol involving three 'all out' sprints of 20s. Despite the interesting preliminary findings regarding the health-related physiological benefits, adherence to these very-low-volume SIT protocols remains unknown. Moreover, it should be noted that these SIT protocols [7,8] were supervised in a laboratory setting. In a 'real world' setting, albeit supervised, Lunt et al. [10] reported drop-out rates were much higher in the SIT and HIIT groups compared to the walking condition (44% vs 18% respectively). Three (19%) SIT participants picked up an injury. These drop-out rates are likely to be much higher in free-living HIIT and SIT programs. It is likely that high degree of effort involved

explained some of the drop-out of participants from the study, which included inactive and overweight participants.

We need not to refer solely to the Dual Mode model (DMM) to describe the negative affective responses to high-intensity exercise. Research has demonstrated a repeatedly negative association between perceived effort and exercise adherence [11]. We recognize that most of this evidence relates to continuous exercise protocols. However, Oliveira et al. [12] found that perceived effort predicted affective response during HIIT and Frazão et al. [13] showed a negative correlation between perceived effort and affective response during HIIT. Wood [14] also found similar declines in affect in SIT and HIIT. In our opinion the negative association between perceived effort and exercise adherence holds for HIIT. Therefore, it is based on this body of evidence and propositions of the DMM that we claim that 'near maximal' and 'supramaximal' interval-training programs are unlikely to produce long-term exercise adherence.

Jung et al. [4] claim that 'interval training' is not that hard and that for inactive/unfit populations it is equated to "walking on a treadmill at a speed of ~3-3.5 mph, at a ~3-5% incline" (~5-6 METs) (p.2) and suggest that relative exercise intensity innate in all prescriptions of 'vigorous' exercise is 'left out'. However, according to the ACSM guidelines [15], ~5-6 METs is equivalent to 'moderate' intensity exercise, and not 'vigorous', for most individuals. Moreover, whilst the speed and incline may not be excessively high, it may be experienced as extremely hard in sedentary populations. Relative exercise intensity refers to the level of physiological stress imposed to subjects' organism (i.e., internal load) [16] and not given walking speed or treadmill inclination (external load markers) as proposed by Jung et al. [17]. Most HIIT protocols use a percentage of peak heart-rate or peak-power output to prescribe the interval work bouts. Thus, less is known about the true relative intensity according to metabolism and its relationship with affective responses and adherence. It seems important that future studies characterize the HIIT protocols according to the ventilatory threshold and respiratory compensation point.

The transfer of SIT and HIIT to an unsupervised setting requiring a high degree of self-regulation and motivation to engage in such 'near maximal' and 'supramaximal' exercise, respectively, is likely to be problematic and further research exploring whether SIT and HIIT can be successfully implemented in a 'real life' setting is necessary. Currently, there is insufficient evidence to promote the efficacy of interval training for public health. Although we agree with Gibala and Hawley [18] that 'SIT is only one option in the armory of primary care interventions that can be used to fight chronic metabolic diseases', we argue that this exercise approach fits for few people and to date the findings from laboratory and supervised studies cannot be transferred to the domain of public health and applied to inactive populations.

In summary, we have argued that the considerable effort required to participate in SIT and HIIT is such that it is likely to damage adherence to such protocols, particularly for independent self-regulation of 'near maximal' and supramaximal efforts protocols. We also argue that SIT and HIIT are unlikely to be experienced as pleasant and enjoyable for most people and that such anticipated displeasure and effort will also damage exercise adherence [19]. Finally, we disagree that disciplinary rivalry exists and note that the previous commentary was co-authored by both physiologists and psychologists working together undertaking HIIT research [13]. We expressed an opinion that such high-intensity training protocols are unlikely to be taken up by most of the sedentary population. We should be united in our pursuit to explore which interventions effectively foster exercise adherence to gain health benefits.

References

1. Hardcastle SJ, Ray H, Beale L, Hagger MS (2014) Why sprint interval training is inappropriate for a largely sedentary population. *Front Psychol* 5:1505.
2. Biddle S, Batterham A (2015) High-intensity interval exercise training for public health: A big HIT or shall we HIT it on the head? *Int J Behav Nutr Phys Act* 12: 95.
3. Weston M, Taylor KL, Batterham AM, Hopkins WG (2014) Effects of low-volume high-intensity interval training on fitness in adults: A meta-analysis of controlled and non-controlled trials. *Sports Med* 44: 1005-1017
4. Jung ME, Little JP, Batterham AM (2016) Commentary: Why sprint interval training is inappropriate for a largely sedentary population. *Front Psychol* 6: 1999.
5. Gillen JB, Gibala MJ (2014) Is high-intensity interval training a time-efficient exercise strategy to improve health and fitness. *Appl Physiol Nutr Metab* 39: 409-412.
6. Saanijoki T, Nummenmaa L, Eskelinen JJ, Savolainen AM, Vahlberg T, et al. (2015) Affective responses to repeated sessions of high-intensity interval training. *Med Sci Sports Exerc* 47: 2604-2611.
7. Metcalfe RS, Babraj JA, Fawcner SG, Niels BJ, Fawcner SG, et al. (2012) Towards the minimal amount of exercise for improving metabolic health: Beneficial effects of reduced-exertion high-intensity interval training. *Eur J Applied Physiol* 112: 2767-2775.
8. Gillen JB, Martin BJ, MacInnis MJ, Skelly LE, Tarnopolsky MA, et al. (2016) Twelve weeks of sprint interval training improves indices of cardiometabolic health like traditional endurance training despite a five-fold lower exercise volume and time commitment. *PLoS ONE* 11: e0154075.
9. Vollard NB, Metcalfe RS (2017) Research into the health benefits of sprint interval training should focus on protocols with fewer and shorter sprints. *Sports Med*.
10. Lunt H, Draper N, Marshall HC, Logan FJ, Hamlin MJ, et al. (2014) High intensity interval training in a real-world setting: a randomized controlled feasibility study in overweight inactive adults, measuring change in maximal oxygen uptake. *PLoS ONE* 9: e83256.
11. Bauman AE, Sallis JF, Dzawaltowski DA, Owen N (2002) Toward a better understanding of the influences on physical activity. The role of determinants, correlates, causal variables, mediators, moderators, and confounders. *Am J Prev Med* 23: 5-14.
12. Oliveira BR, Viana BF, Pires FO, Oliveira M, Santos TM (2015) Prediction of affective responses in aerobic exercise sessions. *CNS Neurol Disord Drug Targets* 14: 1214-1218.
13. Frazão DT, De Farias Junior LF, Dantas TCB, Krinski K, Elsangedy HM, et al. (2016) Feeling of pleasure to high-intensity interval exercise is dependent of the number of work bouts and physical activity status. *PLoS ONE* 11: e0152752.
14. Wood KM, Olive B, LaValle K, Thompson H, Greer K, et al. (2016) Dissimilar physiological and perceptual responses between sprint interval training and high-intensity interval training. *J Strength Cond Res* 30: 244-250.
15. Garber CE, Blissmer B, Deschenes MR, Franklin B, Lamonte MJ, et al. (2011) American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardio-respiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for pre-scribing exercise. *Med Sci Sport Exerc* 43: 1334-1359.
16. Impellizzeri FM, Rampinini E, Marcora SM (2005) Physiological assessment of aerobic training in soccer. *J Sports Sci* 23: 583-592.
17. Jung ME, Bourne JE, Beauvhamp MR, Robinson E, Little JP (2015) High-intensity training as an efficacious alternative to moderate-intensity continuous training for adults with prediabetes. *J Dia Res* 15.
18. Gibala MJ, Hawley JA (2017) Sprinting toward fitness. *Cell Metabolism* 25: 988-990.
19. Zenko Z, Ekkekakis P, Arieli D (2016) Can you have your vigorous exercise and enjoy it too? Ramping intensity down increases post-exercise, remembered, and forecasted pleasure. *J Sport Exerc Psychol* 38: 149-159.